

## Ethnobotanical and Ethnomedicinal Significance of *Myrtus communis* L. in Lower Swat, Pakistan

Qadar Khan<sup>1\*</sup>, Hassan Sher<sup>1</sup>, Jamil Ahmad<sup>1</sup>, Salman Khan<sup>1</sup>, Shamal Habib<sup>1</sup>, Salahud Din<sup>1</sup>, Zarina Rehman<sup>1</sup> and Shazia Dilbar<sup>1</sup>

1. Department of Botany, University of Swat, Khyber Pakhtunkhwa, Pakistan

\*Correspondence: [qadarkhan1131@gmail.com](mailto:qadarkhan1131@gmail.com)

### SUMMARY

*Myrtus communis* L. belongs to the family Myrtaceae and is commonly found in the Lower District, Swat. A survey reported that *Myrtus communis* is traditionally used for medicinal and cultural purposes. Residents provided information about local uses, recipe preparation, and diseases treated, which were recorded. The primary aim of the study was to understand the cultural and ethnomedicinal uses of this species in the study area. Data were collected from 311 informants across eleven localities in Lower Swat: Islamphur (Balavo), Shagai (Ghwara Khela Batorda), Shagai (Shne Shah), Marghazar (Daba Tangai), Marghazar (Manrvano Dub), Barikot (Natmera), Barikot (Nawagai), Barikot (Qwanj), Amlokdara (Gorykan), Jambeel (Shagai), Jambeel (Jhelkanai), and Kokarai (Lokmera/Chamanr). The overarching goal was to involve all participants in gathering information on the medicinal uses and importance of *Myrtus communis*. It is noted that various parts of this species (i.e., leaves, fruits, flowers, stem, and whole plant) are used in cultural practices and to cure different diseases. Culturally, this species is used for tea (n=49), as a flavoring agent (n=17), and as food (n=17). Ethnopharmacologically, it is used to cure chest disorders (n=39), headaches (n=32), hypertension (n=31), diabetes (n=28), stomach issues (n=25), gastric ulcers (n=22), diarrhea (n=16), and tooth pain (n=15). The highest Relative Popularity Level (RPL), i.e., 1.00, is noted for the following cultural uses and diseases: diabetes, stomach issues, gastric ulcers, headaches, chest disorders, hypertension, and tea. The lowest RPL (i.e., 0.14) was noted for anthelmintic and anxiety. The highest Rank Order Priority (ROP) is noted for tea (ROP=15.8), while the ROP for other cultural uses is food (ROP=4.68) and flavoring agents (ROP=4.49).

**Keywords:** *Myrtus communis*, Traditional uses, Swat, Pakistan

**Citation:** Khan, Q., H. Sher, J. Ahmad, S. Khan, S. Habib, S. Din, Z. Rehman, and S. Dilbar. 2026. Ethnobotanical and Ethnomedicinal Significance of *Myrtus communis* L. in Lower Swat, Pakistan. Journal of Wildlife and Ecology. 10(3): 159-170.

Received: 11 April, 2026

Revised: 19 June, 2026

Accepted: 28 July, 2026

Published: 30 September, 2026

### INTRODUCTION

There are about 5,500 species of woody flowering plants in the Myrtaceae family, which is divided into 144 genera and 17 tribes. With 51 genera and roughly 2,500 species mostly found in the Neotropics, the tribe Myrteae accounts for half of the biodiversity of the Myrtaceae family. However, 15 genera and roughly 450 species are found on other continents, including Southeast Asia, Northeast Australia, and the Pacific islands, including New Zealand and New Caledonia. The genus *Myrtus* is the only one found in

Africa and Asia, and Europe (Sisay and Gashaw, 2017; Aboualgasem et al., 2022; Al-Snafi et al., 2024).

*Myrtus communis* develops into a small tree or an evergreen shrub, growing up to 2.5 meters tall. It has a branching stem with dark, evergreen leaves that are glabrous, opposite, glossy, whorled or paired, coriaceous, stiff, lanceolate to oblong, entire-margined, aromatic, acuminate, and 2.5–3.8 cm long. The leaves also lack lamina glands. The axillary flowers, which have a thin peduncle 2 cm in diameter, are starry, fragrant, and can be either pink or white. The stamens have yellow anthers. The blooms, which have five sepals, five petals, and numerous stamens, appear from June to September. The berry fruits, on the other hand, are edible, small, round, and packed with seeds. They ripen in the autumn, from October to February, and are often blue-black, although some varieties have white-yellow fruits. Folk medicine uses several parts of *Myrtus communis* L., including leaves, fruits, flowers, roots, and volatile oils (Dastagir et al., 2022; Giuliani et al., 2022; Al-Snafi et al., 2024). *M. communis* L. is in Sawat known as Manhro.

The habitat of *Myrtus communis* L., a common shrub in the Mediterranean, Asia, and northern subtropical Africa, may have been lost or fragmented as a result of urbanization and agricultural growth (Dabbaghi et al., 2023; Gwynne-Evans et al., 2024). Wild subpopulations are under pressure from increased urbanization, natural fires, livestock grazing, other wild herbivores, and wood cutting. Myrtle fruit and leaf material are frequently gathered from wild subpopulations. However, because it produces many seeds with a high germination rate, the species has high ecological tolerance and is an effective successional species in abandoned fields (Nedjimi, 2024). In general, ethnobotany is the study of human-plant interactions. Some might characterize it as the scientific study of how human cultures interact with plants. The study of plants used for a range of economic and non-commercial societal reasons (e.g., tools and construction materials, food, ritual, divination, cosmetics, adornment, textiles or clothes, cash, and in social life) is included in the purview of ethnobotany (Matsehorova and Odyntsova, 2024).

In Pakistan, the majority of plant studies have focused on their medicinal and other uses (Khan et al., 2017; Umair et al., 2017; Umair et al., 2019; Amjad et al., 2020; Amjad et al., 2021; Hassan et al., 2021; Haidar et al., 2026). While beneficial, these studies often lack information on the plants' significance to people and their various applications. Consequently, some plants may not receive the recognition they merit. *Myrtus communis* L., a significant plant with great cultural and medicinal importance in Lower Swat, is one such example. Little is known regarding its usage in this region. Therefore, *Myrtus communis* L. requires careful research to understand how people utilize it, why it matters to them, and how we can contribute to its protection. This will ensure sustainable use and help preserve local knowledge. *Myrtus communis* L. is a plant that needs our care. By studying it, we can learn more about its significance and how to keep it safe.

## MATERIALS AND METHODS

### STUDY AREA

In district Swat, located in the northern region of Pakistan, the latitudes range from 34° to 35° 55' north and the longitudes from 72° 08' to 72° 50' east (Figure 1). The northern areas include Chitral and Ghizer, to the east are Kohistan and Shangla, while Buner and District Malakand are to the south (Shah *et al.* 2018). The total area of Swat is 5,337 km<sup>2</sup>, with a population of 1,250,000 and a density of 236 residents per km<sup>2</sup>. The elevation in Swat ranges from 760 to 2,280 m. The temperature typically ranges between 22°C and 28°C, with an average rainfall of 864 mm. Most people have one or two cows, from which they sell milk, and they may keep small livestock like goats, sheep, or domesticated birds. Most people are involved in horticulture, using fresh vegetables and fruits, which they also sell (Ali, 2023; Ali et al., 2026).




Figure 1: Map of the study area.

### Study Sites

An ethnobotanical study was conducted from November 2019 to February 2021 to investigate the local uses of medicinal plants in Tehsil Babozai and Barikot, District Swat, Pakistan, in the Hindu Kush mountain range. The selected localities include Islampur (Balavo), Shagai (Ghwara Khela Batorda), Shagai (Shne Shah), Marghazar (Daba Tangai), Marghazar (Manrvano Dub), Barikot (Natmera), Barikot (Nawagai), Barikot (Qwanj), Amlokdara (Gorykan), Jambeel (Shagai), Jambeel (Jhelkanai), and Kokarai (Lokmera/Chamanr), chosen based on the medicinal importance of *Myrtus communis*.

### METHODOLOGY

Data were collected through interviews with 311 respondents, and data on cultural and ethnopharmacological applications were gathered.

## ANALYSIS OF QUANTITATIVE ETHNO-MEDICAL DATA

The data were evaluated using four different indices: frequency of citation (FC), fidelity level (FL), relative popularity level (RPL), and rank order priority (ROP).

### ***Frequency of Citation (FC)***

The frequency of citation (FC) was determined using the total number of informants from the study region (Bashir et al., 2023; Haidar et al., 2026; Jahangeer et al., 2026) who reported the cultural and therapeutic use of *Myrtus communis* during the investigation. Regardless of the number of cultural uses and especially diseases of *Myrtus communis* reported, each informant was counted only once. Therefore, the FC represents the level of recognition and cultural and medicinal use of *Myrtus communis* among the indigenous communities of the study region, with higher FC values indicating greater relevance and cultural significance of the plant.

### ***Fidelity Level (FL)***

The percentage of informants in the study area who identified *Myrtus communis* for a particular medical ailment was determined using the Fidelity Level (FL). The formula is used to compute the FL (Naz et al., 2022):

$$FL (\%) = N_p / FC \times 100$$

Where,

$N_p$  is the number of people who stated *Myrtus communis* for a specific condition.

### ***Relative Popularity Level (RPL)***

The relative popularity of several cultural uses and especially diseases within the study area is shown by the Relative Popularity Index (RPL). It has been noted that the proportion of maximal FC is higher in well-known cultural uses and especially diseases. The remaining cultural uses and especially diseases uses were deemed undesirable. For cultural uses and especially diseases, the relative popularity level was one, but for unpopular cultural uses diseases, it was lower than one (Friedman et al., 1986; Ali-Shtayeh et al., 2000).

### ***Rank Order Priority (ROP)***

Based on their FL and RPL values for the adjustment criterion, the ROP index classifies cultural uses, particularly illnesses (Faiz et al., 2022; Hassan et al., 2022; Rahman et al., 2022). The following formula was used to determine a rank order priority for cultural uses, particularly diseases (Friedman et al., 1986; Ali-Shtayeh et al., 2000):

$$ROP = FL \times RPL$$


### ***Graphical Analysis***

The graphs were designed with the help of MS Excel (2013) and Past statistical software (5.3).

## RESULTS

### PROFILE OF RESPONDENTS


A total of 311 questionnaires were filled out by respondents, of which 244 were male and the remaining 67 were female. They have different occupations, including Teachers (n=24), Students (n=42), Housewives (n=53), Private Employees (n=31), Farmers (n=71), Traditional Healers (n=22), Government Employees (n=48), and Doctors (n=20). They belong to different age groups: 18 to 42 (n=65), 43 to 62 (n=55), 63 to 72 (n=82), and above 72 (n=109). It is noted that most of the respondents (71%) are educated: PhD (n=2), Master's (n=18), Graduate (n=35), Intermediate (n=43), Matric (n=52), and Primary (n=71), while the remaining 29% are illiterate. Most of the respondents belong to rural areas (n=230), while others belong to urban areas (n=81) (Figure 2).



**Figure 2: Profile of respondents of the study area.**

### ATTITUDE OF RESPONDENTS


The reported applications of *Myrtus communis* appeared in all respondents within the given distribution, indicating that its importance was not confined to just one level of respondent reference. The FC above 72 age group showed the greatest overall presence, featuring particularly high rates for different applications, whereas the respondents of 28–42 age group made a significant contribution to multiple reported uses. *Myrtus communis* is widely acknowledged and significant to the Himalayan region people, evident from its presence in both the intermediate and higher age group categories. The category with the most citations indicates that *Myrtus communis* is widely recognized by the surveyed population. In general, the distribution reinforces *Myrtus communis*'s status as an important ethnomedical plant in the study area and indicates a strong level of informant acknowledgment of the plant (Figure 3).



**Figure 3: Ratios of different age group preferences regarding ethnomedicinal uses in the study area.**

#### QAUNTITAVE ANALYSIS

It is noted that various parts of this species (i.e., leaves, fruits, flowers, stem, and whole plant) are used in cultural practices and to cure different diseases (Figure 4).




**Figure 4: Uses of parts of *Myrtus communis* L. from study area.**


#### Frequency of Citation (FC)

It is noted that, culturally, this species is used for tea (n=49), flavoring agent (n=17) and food (n=17); while ethnopharmacologically, it is used to cure chest disorders (n=39), headaches (n=32), hypertension (n=31), diabetes (n=28), stomach issues (n=25), gastric ulcers (n=22), as a diarrhea (n=16), and tooth pain (n=15) (Figure 5). It is noted that a higher tendency of uses (n=122) of this species was found in the oldest

age group (above 70), followed by the age group 28-42 (n=79), the age group 63-72 (n=60), and the age group 43-62 (n=50) (Figure 6).



**Figure 5: Cultural and traditional medicinal Uses of species in the study area.**



**Figure 6: Trends in species usage across different age groups.**

### ***Fidelity Level (FL)***


The species with the highest therapeutic value, which corresponds to the highest FL, has been discovered in the research region. Tea (FL=15.80) has the highest FL value, followed by Chest disorder (FL=12.54), Headache (FL=10.29), Hypertension (FL=9.97), Diabetes (FL=9.00), Stomach (FL=8.04), Gastric ulcer (FL=7.07), Food (FL=5.70), Flavoring agent (FL=5.47), Diarrhea (FL=5.14), Teeth pain (FL=4.82), Abdominal pain (FL=2.89), Fever (FL=1.61), and Anthelmintic (FL=0.96). By contrast, Anxiety (FL=0.96) has the lowest percentage, which was seen in Table 1 and Figure 7.

### ***Relative Popularity Level (RPL)***

The highest RPL, i.e., 1.00, is noted for the following cultural uses and especially diseases: Diabetes, Stomach, Gastric ulcer, Headache, Chest disorder, Hypertension, and Tea; while the lowest RPL (i.e., 0.14) was noted for Anthelmintic and Anxiety. Additionally, RPL was noted for other cultural uses as follows: Flavoring agent (RPL=0.82), Food (RPL=0.82), Diarrhea (RPL=0.77), Teeth pain (RPL=0.72), Abdominal pain (RPL=0.43), and Fever (RPL=0.24) (Table 1 and Figure 7).

### ***Rank Order Priority (ROP)***

The highest Rank Order Priority is noted for Tea (ROP=15.8), followed by Chest Disorder (ROP=12.5), Headache (ROP=10.3), Hypertension (ROP=9.97), Diabetes (ROP=9), Stomach (ROP=8.038585), Gastric Ulcer (ROP=7.07), Diarrhea (ROP=3.98), Tooth Pain (ROP=3.495), Abdominal Pain (ROP=1.26), Fever (ROP=0.388), Anthelmintic (ROP=0.1398), and Anxiety (ROP=0.1398) for diseases, while the Rank Order Priority for cultural uses is Food (ROP=4.68) and Flavoring Agent (ROP=4.489) (Table 1 and Figure 7).



**Figure 7: Comparison of FL, RPL and ROP.**

**Table 1: Cultural and medicinal and daily uses of *Myrtus communis* in Lower Swat, Pakistan.**

| Uses            | FC<br>(28-42) | FC<br>(43-62) | FC<br>(63-72) | FC<br>(above 72) | FC | FL    | RPL  | ROP   |
|-----------------|---------------|---------------|---------------|------------------|----|-------|------|-------|
| Diabetes        | 1             | 3             | 6             | 18               | 28 | 9.00  | 1.00 | 9.00  |
| Stomach         | 11            | 0             | 8             | 6                | 25 | 8.04  | 1.00 | 8.04  |
| Anthelmintic    | 2             | 0             | 0             | 1                | 3  | 0.96  | 0.14 | 0.14  |
| Gastric ulcer   | 10            | 3             | 2             | 7                | 22 | 7.07  | 1.00 | 7.07  |
| Headache        | 4             | 8             | 3             | 17               | 32 | 10.29 | 1.00 | 10.29 |
| Fever           | 2             | 0             | 1             | 2                | 5  | 1.61  | 0.24 | 0.39  |
| Abdominal pain  | 0             | 7             | 2             | 0                | 9  | 2.89  | 0.43 | 1.26  |
| Chest disorder  | 12            | 4             | 8             | 15               | 39 | 12.54 | 1.00 | 12.54 |
| Anxiety         | 1             | 0             | 0             | 2                | 3  | 0.96  | 0.14 | 0.14  |
| Diarrhea        | 0             | 2             | 5             | 9                | 16 | 5.14  | 0.77 | 3.98  |
| Hypertension    | 1             | 5             | 8             | 17               | 31 | 9.97  | 1.00 | 9.97  |
| Teeth pain      | 9             | 2             | 0             | 4                | 15 | 4.82  | 0.72 | 3.50  |
| Flavoring agent | 5             | 3             | 7             | 2                | 17 | 5.47  | 0.82 | 4.49  |
| Food            | 2             | 9             | 5             | 1                | 17 | 5.70  | 0.82 | 4.68  |
| Tea             | 19            | 4             | 5             | 21               | 49 | 15.80 | 1.00 | 15.80 |

## DISCUSSION

In the present study, it is noted that leaves, fruits, flowers, stems, and the whole plant are used culturally (i.e., as flavoring agents, food, and tea) and to cure different diseases (i.e., diabetes, stomach issues, anthelmintic conditions, gastric ulcers, headaches, fever, abdominal pain, chest disorders, anxiety, diarrhea, hypertension, and tooth pain). In the past, *Myrtus communis* was utilized in traditional medicine, herbal remedies, and a variety of cultural practices (Rezaee et al., 2025). Large numbers of people in the modern world who lack access to modern medical facilities also utilize myrtle. Herbal medicines differ from synthetic treatments due to their safety profile, which includes minimizing adverse effects, cost-effectiveness, and natural healing qualities (Dabbaghi et al., 2023). In the Mediterranean, Middle Eastern, and South Asian regions, myrtle has long been used as traditional medicine. Pakistan, Iran, India, Turkey, and Greece are the most prominent of these (Ullah et al., 2025). Gastrointestinal disorders, skin conditions, renal and urinary disorders, musculoskeletal disorders, respiratory infections, eye infections, oral infections, parasitic infections, gynecological conditions, ENT-related issues, fever, cancer, and other ailments are among the various diseases that myrtle has been reported to treat. The cumulative percentage of gastrointestinal disorders that myrtle heals is due to its medicinal properties, as well as its digestive stimulant and antimicrobial effects (Rezaee et al., 2025).

Additionally, traditional myrtle healers reported using the leaves and essential oil to cure cuts, wounds, burns, boils, dermatitis, and fungal infections. These have astringent and antibacterial properties that aid in preventing infection during aftercare. Myrtle extracts have been detected in ointments and herbal pastes for wound care. In addition, it possesses antifungal properties against common skin conditions like fungal dermatitis and ringworm. Additionally, myrtle is utilized to maintain renal and urinary tract health. The literature most frequently mentions the leaves and roots as being useful in treating certain conditions. Physicians employed myrtle extracts to make remedies

for bladder problems, kidney stones, and UTIs. Additionally, a variety of phenolic compounds found in plant parts aid in kidney stone dissolution and enhance renal function in general (Rezaee et al., 2025; Aldosary et al., 2026; Tahralli, 2026).

Over the past 20 years, there has been significant growth in demand for herbal medicines. To ensure the safety, effectiveness, and consistency of herbal medicines, strict quality controls have been implemented. Phytochemical evaluation is the best method for assessing the quality of herbal medication. These methods can also be used to standardize herbal formulations for therapeutic use and identify the classes of bioactive chemicals that give herbs their medicinal qualities (Matsehorova and Odyntsova, 2024). Flavonoids, tannins, terpenoids, alkaloids, phenolic acids, and essential oils are just a few of the phytochemicals found in myrtle. Berries and leaves are known to contain flavonoids and tannins, which have potent antibacterial, anti-inflammatory, and antioxidant properties. The essential oil extract from the myrtle plant is primarily composed of terpenoids, which give it its distinctive scent and antibacterial properties (Yarahmadi et al., 2024). Nonetheless, there are many bioactive substances in the phytochemical composition of myrtle. Twelve of the sixty compounds found in the myrtle plant are volatile organic compounds, including phenols, alcohols, terpenoids, fatty acids, sterols, glycosides, carboxylic acids, flavonoids, derivatives of cinnamic acid, and nitro compounds. These chemicals are responsible for the plant's pharmacological activity, including its antibacterial, antioxidant, and anti-inflammatory properties. In terms of the bioactive compounds utilized in both herbal therapy and contemporary pharmaceutical medications, myrtle plays a significant role (Aboud et al., 2022; Al-Snafi et al., 2024).

## CONCLUSIONS

*Myrtus communis* medicinally very important and is used by the local people of Lower Swat, Pakistan. *M. communis* L. is known in Swat as Manhro. The leaves, fruits, flowers, stem, and whole plant are used for tea (n=49), flavoring agents (n=17), and food (n=17); while ethnopharmacologically, it is used to cure chest disorders (n=39), headaches (n=32), hypertension (n=31), diabetes (n=28), stomach issues (n=25), gastric ulcers (n=22), diarrhea (n=16), and tooth pain (n=15). Due to this information, we collect data, but *Myrtus communis* requires special attention and support to conserve and ensure safety, so that poor people can easily access medication in the future.

## REFERENCES

- Aboualgasem, F. A., H. M. Abozweta, A. E. Mohamed, and A. S. Elghdafi. 2022. *Myrtus communis* L. phytochemistry, ethnobotany and pharmacology: A review. The Libyan Association. 1-21.
- Aboud, M. N., A. D. M. Al-Maliki, A. M. Abduljeel, and M. F. Hameed. 2022. Estimation and Evaluation of Antibacterial Efficacy of Phenolics Isolated from Iraqi *Myrtus communis* Leaves. Teikyo Medical Journal. 45: 6853-6859.
- Al-Snafi, A. E., J. O. Teibo, H. M. Shaheen, O. A. Akinfe, T. K. A. Teibo, N. Emieseimokumo, M. M. Elfiky, H. M. Al-Kuraishy, A. I. Al-Garbeeb, and A. Alexiou. 2024. The therapeutic value of *Myrtus communis* L.: an updated review. Naunyn-Schmiedeberg's archives of pharmacology. 397: 4579-4600.
- Aldosary, N., I. Alnajim, M. Al-Darwish, R. Alasadi, H. Ali, and F. Al-Hamoud. 2026. Evaluation of the effect of nanoemulsion formulation of *Myrtus communis* essential oil against date palm white scale insect *Parlatoria blanchardi*. Plant Protection (Scientific Journal of Agriculture). 49: 23-40.
- Ali-Shtayeh, M. S., Z. Yaniv, and J. Mahajna. 2000. Ethnobotanical survey in the Palestinian area: a classification of the healing potential of medicinal plants. Journal of ethnopharmacology. 73: 221-232.
- Ali, S. 2023. A study on different plant species of the Rosaceae family and their ethnobotanical uses among the local communities at Swat district, Pakistan. Ethnobotany Research and Applications. 25: 1-16.

- Ali, S., S. A. Shah, and S. Din. 2026. An ethnobotanical study of wild plant resources used among locals of high mountainous regions of Swat Valley, Pakistan. *Ethnobotany Research and Applications*. 34: 1-37.
- Amjad, M. S., R. Khurshid, A. M. Abbasi, M. Altaf, H. Qureshi, and F. Abasi. 2021. Ethnobotany of Western Himalayan Region, Azad Jammu and Kashmir, Pakistan. *Ethnobiology of Mountain Communities in Asia*. 87.
- Amjad, M. S., U. Zahoor, R. W. Bussmann, M. Altaf, S. M. H. Gardazi, and A. M. Abbasi. 2020. Ethnobotanical survey of the medicinal flora of Harigal, Azad Jammu & Kashmir, Pakistan. *Journal of Ethnobiology and Ethnomedicine*. 16: 65.
- Bashir, S., M. Altaf, T. Hussain, M. Umair, M. Majeed, W. Mangrio, A. Khan, A. Gulshan, M. Hamed, and S. Ashraf. 2023. Vernacular Taxonomy, Cultural and Ethnopharmacological Applications of Avian and Mammalian Species in the Vicinity of Ayubia National Park, Himalayan Region. *Biology* 12: 609.
- Dabbaghi, M. M., M. S. Fadaei, H. Soleimani Roudi, V. Baradaran Rahimi, and V. R. Askari. 2023. A review of the biological effects of *Myrtus communis*. *Physiological reports*. 11: e15770.
- Dastagir, G., I. Ahmad, and N. U. Uza. 2022. Micromorphological evaluation of *Daphne mucronata* Royle and *Myrtus communis* L. using scanning electron microscopic techniques. *Microscopy Research and Technique*. 85: 1120-1134.
- Faiz, M., M. Altaf, M. Umair, K. S. Almarry, Y. B. Elbadawi, and A. M. Abbasi. 2022. Traditional uses of animals in the Himalayan region of Azad Jammu and Kashmir. *Frontiers in Pharmacology*. 13: 807831.
- Friedman, J., Z. Yaniv, A. Dafni, and D. Palewitch. 1986. A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev Desert, Israel. *Journal of Ethnopharmacology*. 16: 275-287.
- Giuliani, C., M. Bottoni, F. Milani, S. Toderò, P. Berera, F. Maggi, L. Santagostini, and G. Fico. 2022. Botanic Garden as a Factory of Molecules: *Myrtus communis* L. subsp. *communis* as a Case Study. *Plants*. 11: 754.
- Gwynne-Evans, D., D. M. Richardson, and R. McKenzie. 2024. *Myrtus communis* (Myrtaceae) as an alien species in South Africa: Status and prognosis. *South African Journal of Botany*. 166: 550-560.
- Haidar, R., M. Altaf, M. Faiz, S. T. u. Shukurova, O. Shukurov, and A. M. Abbasi. 2026. Ethnomedicinal uses of fauna in the Lesser Himalayan Region of Pakistan. *Ethnobotany Research and Applications*. 34: 1-23.
- Hassan, M., S. M. Haq, M. Majeed, M. Umair, H. A. Sahito, M. Shirani, M. Waheed, R. Aziz, R. Ahmad, and R. W. Bussmann. 2022. Traditional food and medicine: Ethno-traditional usage of fish Fauna across the valley of Kashmir: A western Himalayan Region. *Diversity*. 14: 455.
- Hassan, M., S. M. Haq, U. Yakoob, M. Altaf, and R. W. Bussmann. 2021. The ethnic diversities in animal-human interactions in former Jammu and Kashmir State-India. *Ethnobotany Research and Applications*. 22: 1-18.
- Jahangeer, M., M. Altaf, N. Irshad, M. Tariq, S. Chowhan, T. Hussain, A. H. Narejo, M. Ali, M. Muhamad, and S. Ashraf. 2026. Cultural and ethnopharmacological uses of wildlife in the himalayan region of Azad Jammu and Kashmir, Pakistan. *Journal of Ethnobiology and Ethnomedicine*. 22(13): 1-35.
- Khan, A., S. Mehmood, and R. A. Khan. 2017. Ethnobotanical study of some wild herb medicinal Xerophytes of district Bannu, Khyber Pakhtunkhwa, Pakistan. *J. Wildl. Ecol.* 1: 37-51.
- Matsehorova, O. Y., and V. Odyntsova. 2024. Prospects for the creation of new phytochemical medicinal products based on *Myrtus communis* L.(a review). ;17(1):70-78.
- Naz, A., M. S. Amjad, M. Umair, M. Altaf, and J. Ni. 2022. Medicinal plants used as therapeutic medicine in the Himalayan region of Azad Jammu and Kashmir. *South African Journal of Botany*. 150: 194-216.
- Nedjimi, B. 2024. Common myrtle (*Myrtus communis* L.): trace element composition and contribution to dietary intake. *Food Chemistry Advances*. 4: 100644.
- Rahman, Q., M. S. Nadeem, M. Umair, M. Altaf, J. Ni, A. M. Abbasi, M. A. Jameel, A. Pieroni, M. H. Hamed, and S. Ashraf. 2022. Medicinal waterbirds in the traditional healthcare system: an assessment of biodiversity-cultural linkages in Eastern Khyber Pakhtunkhwa, Pakistan. *Journal of ethnobiology and ethnomedicine*. 18: 1-19.
- Rezaee, A., M. Rezaee, Z. Feyzabadi, and M. M. Zarshenas. 2025. Clinical Applications of *Myrtus communis* L. in Traditional and Modern Medicine: A Scoping Review. *Pharmaceutical & Biomedical Research*. 11: 173.
- Sisay, M., and T. Gashaw. 2017. Ethnobotanical, ethnopharmacological, and phytochemical studies of *Myrtus communis* Linn: A popular herb in Unani system of medicine. *Journal of evidence-based complementary & alternative medicine*. 22: 1035-1043.
- Tahralli, I. 2026. Anti-inflammatory effects of *Myrtus communis* L. (myrtle): experimental and clinical evidence with an immunological perspective: Ilhan Tahralli. *Inflammopharmacology*. 34: 2175-2192.
- Ullah, F., T. Jamal, W. Ahmad, M. Usama, A. Ullah, K. Khan, S. Jabeen, and A. Basir. 2025. *Myrtus communis* L. (Myrtle): A comprehensive Review of its Folk uses, Medicinal uses, Historical perspective, Phytochemistry, and Pharmacology. *Journal of Agriculture Sustainability and Environment*. 4: 38-56.

- Umair, M., M. Altaf, and A. M. Abbasi. 2017. An ethnobotanical survey of indigenous medicinal plants in Hafizabad district, Punjab-Pakistan. PloS one. 12: e0177912.
- Umair, M., M. Altaf, R. W. Bussmann, and A. M. Abbasi. 2019. Ethnomedicinal uses of the local flora in Chenab riverine area, Punjab province Pakistan. Journal of Ethnobiology and Ethnomedicine. 15(7):1-21.
- Yarahmadi, R., H. Mumivand, A. Ehtesham Nia, M. R. Raji, and S. Argento. 2024. Natural diversity in total phenol, flavonoids, antioxidant properties, and essential oil composition of Iranian populations of *Myrtus communis* L. Plants. 13: 3458.

**Competing interests:** Authors have declared that no competing interests exist.

**Funding:** Authors have no source of funding for this work.

**Authors' contributions:** Data were collected and a manuscript was written by Khan; statistically analyzed by Khan; and critically reviewed and approved as final by all authors.

